

The formation and retention of gas giant planets around stars with a range of metallicities

Shigeru Ida

Tokyo Institute of Technology, Ookayama, Meguro-ku, Tokyo 152-8551, Japan

ida@geo.titech.ac.jp

and

D. N. C. Lin

UCO/Lick Observatory, University of California, Santa Cruz, CA 95064

lin@ucolick.org

ABSTRACT

The apparent dependence of detection frequency of extrasolar planets on the metallicity of their host stars is investigated with Monte Carlo simulations using a deterministic core-accretion planet formation model. According to this model, gas giants formed and acquired their mass M_p through planetesimal coagulation followed by the emergence of cores onto which gas is accreted. These protoplanets migrate and attain their asymptotic semi-major axis a through their tidal interaction with their nascent disk. Based on the observed properties of protostellar disks, we generate M_p - a distribution. Our results reproduce the observed lack of planets with intermediate mass $M_p = 10$ – $100M_\oplus$ and $a \lesssim 3\text{AU}$ and with large mass $M_p \gtrsim 10^3 M_\oplus$ and $a \lesssim 0.2\text{AU}$. Based on the simulated M_p - a distributions, we also evaluate the metallicity dependence of fraction of stars harboring planets that are detectable with current radial velocity survey. If protostellar disks attain the same fraction of heavy elements which are contained in their host stars, the detection probability around metal-rich stars would be greatly enhanced because protoplanetary cores formed in them can grow to several Earth masses prior to their depletion. These large masses are required for the cores to initiate rapid gas accretion and to transform into giant planets. The theoretically extrapolated metallicity dependence is consistent with the observation. This correlation does not arise naturally in the gravitational-instability scenario. We also suggest other metallicity dependence of the planet distributions that can be tested by on-going observations.

Subject headings: planetary systems: formation – solar system: formation – stars: statics

1. Introduction

One of the most important observed characteristics of known extrasolar planets is that their detection frequency η_J increases with the metallicity Z_* of their host stars (Fischer & Valenti 2003; Santos et al. 2004). Although Z_* could be significantly changed by the accretion of planets/planetesimals onto the host stars after they have evolved onto the main sequence, e.g., (Sandquist et al. 1998; Murray & Chaboyer 2002), the lack of any Z_* dispersion among members of stellar clusters (Quillen 2002; Wilden et al. 2002) indicate that the impact of this process is very limited. The η_J - Z_* correlation may also be interpreted as evidence that the formation probability of gas giant planets is greatly enhanced in metal-rich protostellar disks, in accordance with the core-accretion scenario for formation of giant planets, e.g., (Mizuno 1980; Bodenheimer & Pollack 1986; Pollack et al. 1996; Ikoma et al. 2000).

In the conventional core-accretion scenario, heavy elements (metals) in a protoplanetary disk condense into grains which form km-sized planetesimals. Through gravitational interaction and coagulation, planetesimals evolve into cores through runaway (Greenberg et al. 1978; Wetherill & Stewart 1989; Aarseth et al. 1993; Kokubo & Ida 1996) and oligarchic growth (Kokubo & Ida 1998, 2002). The cores’ eccentricity is effectively damped by their tidal interaction with the ambient disk gas (Artymowicz 1993; Ward 1993). After the cores have swept up all residual planetesimals within a “feeding-zone width” ~ 10 Hill’s radii (Kokubo & Ida 1998, 2002), their growth in the inner regions of the disks is stalled with an “isolation” mass $M_{c,iso}$. If a core attains the critical mass, $M_{c,acc} \sim \text{several } M_{\oplus}$, before the disk gas depletion, rapid gas accretion onto it is initiated and it transforms into a gas giant planet. Since $M_{c,iso}$ increases with the surface density (Σ_d) of dusts (composed of metals) in protoplanetary disks, formation of gas giants tends to be more prolific in a metal-rich environment.

Here, using the deterministic planet formation model developed by Ida & Lin (2004) (hereafter referred to as Paper I), we calculate the metallicity dependence of fraction (η_J) of stars harboring planets that are detectable with the current radial velocity surveys. In §2, we briefly describe our model. In §3, we simulate the mass (M_p) – semimajor axis (a) distributions of extrasolar planets and their metallicity dependence through a series of Monte Carlo simulations. The theoretically constructed distributions are consistent with observed ones. Using the simulated distributions, we evaluate η_J . Our model explains the observed metallicity dependence of η_J . §4 is summary and discussions.

2. Model

We carry out Monte Carlo simulations to reproduce the observed M_p - a distribution of extrasolar planets. We basically follow the methods of Paper I except slightly different choice of parameters. Initial conditions of the Monte Carlo simulations are the a of planets and the dust and gas surface density (Σ_g and Σ_d) at a in their nascent disks. For a given set of $\Sigma_g(a)$ and $\Sigma_d(a)$,

we numerically compute the entire formation and migration sequence of individual cores and giant planets. Here we briefly summarize our prescription. For details, see Paper I.

Since their observational data over the relevant length scales (a few AU’s) for planet-forming regions is not available, we scale their quantitative values with global factors f_d and f_g , to those of the empirical minimum mass nebula (MMN) model for our Solar system as

$$\begin{cases} \Sigma_d &= 10\eta_{\text{ice}}f_d(a/1\text{AU})^{-3/2} \text{ [g cm}^{-2}\text{]}, \\ \Sigma_g &= 2.4 \times 10^3 f_g(a/1\text{AU})^{-3/2} \text{ [g cm}^{-2}\text{]}, \end{cases} \quad (1)$$

where the compositional scaling parameter $\eta_{\text{ice}} = 1$ inside the ice condensation radius [$a_{\text{ice}} = 2.7(M_*/M_\odot)^2$ AU and 4.2 outside it (Hayashi 1981). [Note that η_{ice} outside a_{ice} can be slightly smaller (~ 3.0) (Pollack et al. 1994).] The mass of the host star M_* is scaled with that of the Sun $M_\odot$. The dependence on the power-law index for Σ_d and Σ_g are examined in a separate paper. To represent the decline in the observational signatures of protostellar disks on timescales $\tau_{\text{dep}} \sim 1\text{--}10$ Myrs (Haisch et al. 2001), we assume exponential decay of Σ_g as

$$f_g = f_{g,0} \exp(-t/\tau_{\text{dep}}), \quad (2)$$

due to both viscous evolution and photoevaporation. We neglect gas replenishment onto the disk and any time variation in f_d . We discuss the distributions of f_d and $f_{g,0}$ below.

With prescription (1), the cores’ mass at time t after the formation of the disk is deduced to be (Paper I)

$$M_c(t) \simeq \left(\frac{\tau_{\text{dep}}}{4.8 \times 10^5 \text{ years}} \right)^3 \eta_{\text{ice}}^3 f_d^3 f_{g,0}^{6/5} \left(\frac{m}{10^{22} \text{ g}} \right)^{-2/5} \left(\frac{a}{1\text{AU}} \right)^{-81/10} \left(\frac{M_*}{M_\odot} \right)^{1/2} \left(1 - \exp \left(\frac{-2t}{5\tau_{\text{dep}}} \right)^3 \right) M_\oplus. \quad (3)$$

The dependence of $M_c(t)$ on the mass m of the field planetesimals is due to the aerodynamical gas drag which regulates their velocity dispersion. Since e-folding growth time scale is longer for larger M_c , $M_c(t)$ does not depend on initial mass of the core. In the nominal case, Eq. (3) with $m = 10^{22}$ g is used. If m is smaller or if radial migration of cores due to their tidal interaction with gas (Goldreich & Tremaine 1980; Ward 1986) and/or planetesimal disks (Murray et al. 1998; Ida et al. 2000) occurs, core accretion may be accelerated (Inaba et al. 2003; Tanaka & Ida 1999; Rice & Armitage 2003). We also carry out the calculations with 3 times larger/smaller dM_c/dt . Note that we do not consider the change in a of the cores due to the radial migration, because the migration can be a random walk (Nelson & Papaloizou 2004).

Prior to severe depletion, the cores’ growth is stalled when they attain an isolation mass (Kokubo & Ida 1998; Kominami & Ida 2002) which is given by

$$M_{c,\text{iso}} \simeq 0.16 \eta_{\text{ice}}^{3/2} f_d^{3/2} \left(\frac{a}{1\text{AU}} \right)^{3/4} \left(\frac{M_*}{M_\odot} \right)^{-1/2} M_\oplus. \quad (4)$$

In our simulation, growth beyond the isolation after significant gas depletion is taken into account (Paper I). If radial migration of the cores occurs, their growth barrier can be bypassed (Tanaka

& Ida 1999; Rice & Armitage 2003) up to $M_{\text{c,noiso}}$, which is given with $M_{\text{c,noiso}} \simeq \pi \Sigma_{\text{d}} a^2$ (feeding zone width $\Delta a \sim a$) by (Paper I)

$$M_{\text{c,noiso}} \simeq 1.2 \eta_{\text{ice}}^{3/2} f_{\text{d}} \left(\frac{a}{1\text{AU}} \right)^{1/2} M_{\oplus}. \quad (5)$$

In the nominal case, the asymptotic masses of the cores are given by $\min(M_{\text{c,iso}}, M_{\text{c,noiso}})$. We also examine the case where $M_{\text{c,noiso}}$ is adopted in entire f_{d} and a .

For sufficiently rapid gas accretion to transform a core into a gas giant planet, it must attain a critical core mass, $M_{\text{c,acc}} \sim \text{several } M_{\oplus}$, before the disk gas depletion. There are at least two factors which limit the gas accretion rate. Planetary atmosphere is no longer in hydrostatic equilibrium and gas accretion starts when M_{c} exceeds $M_{\text{c,hydro}} \simeq 10(\dot{M}_{\text{c}}/10^{-6} M_{\oplus} \text{yr}^{-1})^{0.25} M_{\oplus}$ (Ikoma et al. 2000). Pollack et al. (1996) suggested that $M_{\text{c,hydro}} > 10 M_{\oplus}$ because $\dot{M}_{\text{c}}$ may be maintained at a modest value, for a few Myr. But, cores emerge with neighbors of comparable masses. Together they stir up the random motion of the residual planetesimals, widen their feeding zone, and reduce their $\dot{M}_{\text{c}}$. In addition, the combined effects of gravitational scattering by the cores and gas drag actually lead to a clearing of residual planetesimals in the feeding zones of the cores (Tanaka & Ida 1997; Rafikov 2003) and a reduction in $M_{\text{c,hydro}}$. Thus, the onset of efficient gas accretion would occur earlier than previously estimated.

With a negligible $\dot{M}_{\text{c}}$, the gas accretion rate is determined by the planet’s total (core plus envelope) mass (M_{p}) such that $dM_{\text{p}}/dt = M_{\text{p}}/\tau_{\text{KH}}$ where We adopt

$$\tau_{\text{KH}} = 10^b \left(\frac{M_{\text{p}}}{M_{\oplus}} \right)^{-c} \text{ yrs}, \quad (6)$$

with $b = 10$ and $c = 3$, which is a fitting formula for Pollack et al. (1996)’s results with negligible planetesimal accretion. τ_{KH} for $M_{\text{p}} \sim 1\text{--}10 M_{\oplus}$ can be longer if modest planetesimal accretion induced by gas accretion is assumed [e.g., “phase 2” by Pollack et al. (1996)]. But, the planetesimal accretion may be significantly reduced as discussed above. In this case, phase 2 cannot be sustained. On the other hand, if the core’s migration is very fast, the growth of a core does not decline until it acquires mass $\sim M_{\text{c,noiso}}$. In this case, phase 2 does not exist either (Alibert et al. 2004). So, in the nominal case, we adopt $(b, c) = (10, 3)$, but $(b, c) = (9, 3)$ and $(11, 3.5)$ are also examined (the latter corresponds to the gas accretion including phase 2).

The gas accretion rate increases rapidly with M_{p} . Such runaway gas accretion is terminated when the residual gas is depleted globally on timescale τ_{dep} or it is severely depleted locally in the vicinity of their orbits due to gap opening, which is assumed to occur if their Hill radius exceeds 1.5 times the disk scale height (Paper I). When M_{p} becomes a significant fraction of M_{J} (Jupiter mass), planets’ tidal torque on the disk becomes larger than the viscous torque of the disk gas, resulting in a partial gap opening, e.g., (Lubow et al. 1999). It reduces gas accretion rate. Bondi accretion limit also becomes important in the reduction. However, the elongated accretion timescale may be still shorter than τ_{KH} for $M_{\text{p}} \sim \text{several } M_{\oplus}$, so that the limited accretion at high M_{p} may not affect

the mass distribution of gas giants significantly. The critical mass $M_{c,acc}$ for actual formation of gas giants corresponds to the value of M_p for which $\tau_{KH} \sim \tau_{dep}$. Equation (6) shows $M_{c,acc} \sim$ several $M_\oplus$. From eqs. (4) and (3), we find that, for sufficiently large f_d , it is possible for $M_c(\tau_{dep}) > M_{c,acc}$ and $M_{c,iso} > M_{c,acc}$ so that gas giant can form readily.

After the gap opening, the planets undergo “type-II” migration, in contrast to the cores’ “type-I” migration without a gap (Ward 1997). In order to compute the protoplanets’ migration, we adopt the α -prescription for effective viscosity (Shakura & Sunyaev 1973) with a uniform $\alpha = 10^{-4}$ for all the disks such that their viscous diffusion time scale (τ_ν) at $a \sim 10$ AU is comparable to $\tau_{dep} = 1$ –10 Myrs. The planets migrate (Lin & Papaloizou 1985, 1993) with a viscously evolving disk on time scales (Paper I)

$$\tau_{mig} = \frac{a}{\dot{a}} = 10^6 f_g^{-1} \left(\frac{M_p}{M_J} \right) \left(\frac{a}{1 \text{ AU}} \right)^{1/2} \text{ yrs.} \quad (7)$$

The planets’ migration is terminated either when the disk is severely depleted or when they reach the vicinity of their host stars. We artificially halt the planetary migration at 0.04 AU.

Using these models, we carry out Monte Carlo simulations. For simplicity, we generate a set of initial a ’s of the protoplanets, τ_{dep} and M_* with uniform distributions in log scale in the ranges of 0.1–100 AU, 10^6 – 10^7 yrs, and 0.7 – $1.4 M_\odot$, respectively. The assumed a distribution corresponds to orbital separations Δa that is proportional to a , which may be the simplest choice. More refined choice of Δa that takes into account interactions between planets and detailed formation process will be explored in a future paper. However, as long as Δa increases with a with a moderate power index, calculated distributions of gas giants may not change significantly.

We discuss the distribution of f_d and $f_{g,0}$ in details, because it is a key assumption of this paper. A large range of dust mass in a disk is inferred. Observationally, the mass transfer rate from disks onto classical T Tauri stars is $\sim 10^{-7}$ – $10^{-9} M_\odot \text{ yr}^{-1}$ (Calvet et al. 2000) with inferred disk masses equivalent to $0.1 \lesssim f_{g,0} \lesssim 10$. Independent mm observations (Beckwith & Sargent 1996) show a similar range of f_d . Here we assume that $\log_{10} f_{g,0}$ has a Gaussian distribution centered at 0 with the dispersion 1.0 and upper cut-off at $f_{g,0} = 30$ (Paper I).

Since in the MMN, $f_d = f_{g,0} = 1$, the independently determined mean values $\langle f_d \rangle \sim \langle f_{g,0} \rangle \sim 1$ imply that the average value of metallicity in the disk (Z_{disk}) is $\sim Z_\odot$ (the Solar composition), where $[Z_{\text{disk}} \equiv \log_{10}(f_d/f_{g,0})]$. In Paper I, $f_d = f_{g,0}$ was assumed. Here we extend our calculation to a range of Z_{disk} in order to account for the most pronounced correlation between the percentage of stars with planets detectable with the current Doppler survey and metallicity of their host stars (Z_*) (Fischer & Valenti 2003; Santos et al. 2004), assuming $Z_* \sim Z_{\text{disk}}$. From this inference, we explore the possibility that $\langle f_d \rangle$ depends on Z_* as $\langle f_d \rangle = 10^{Z_{\text{disk}} - Z_\odot} \sim 10^{Z_* - Z_\odot}$, while $\langle f_{g,0} \rangle$ is ~ 1 , independent of Z_* , since gas disk formation/evolution processes may not be significantly affected by Z_{disk} ($\sim Z_*$). We assume $f_d = f_{g,0} \times 10^{Z_* - Z_\odot}$. This estimate can be verified with additional direct measurement of both gas and dust masses in protostellar disks in the near future. All stellar material is accreted through protostellar disks where differential retention of gas and dust would affect the stellar metallicity. These key assumptions are supported by the remarkable chemical

homogeneity among young stars in the Pleiades cluster (Wilden et al. 2002) and mature binary systems (Desidera et al. 2004).

3. Numerical Results

The simulated M_p – a distributions with $\Delta Z_* \equiv Z_* - Z_\odot = 0$ are shown in Figure 1a. This is the nominal case where $m = 10^{22}$ g, $M_{\text{c,iso}}$ and $\tau_{\text{KH}} = 10^{10}(M_p/M_\oplus)^{-3}$ yrs are adopted. For comparison, the observed distributions are also plotted in Figure 1b. The observational limits in the current Doppler survey, $M_p \gtrsim 100(a/1\text{AU})^{1/2}M_\oplus$ and $a \lesssim 3\text{AU}$ (around F, G, K dwarfs within ~ 50 pc) are marked by dotted lines. A deficit of planets with an asymptotic mass $M_p = 10\text{--}100M_\oplus$ and $a \lesssim 3\text{AU}$ (within a “planet desert”) is apparent in Fig. 1a. This sparsely populated region is primarily caused by the runaway gas accretion onto cores with $M_p \gtrsim$ several $M_\oplus$ (Paper I). This region divides the terrestrial (rocky) planet, gas and ice giant domains in the planets’ mass-period distribution. Planetary migration generally sharpens the boundaries of the domains (Paper I; Udry et al. (2003)). Since migration is slower for larger M_p (Eq. [7]), relatively massive planets are less likely to migrate to inner regions, resulting in a deficit of planets with $M_p \gtrsim 10^3M_\oplus$ at $a \lesssim 0.2\text{AU}$ (Fig. 1a; also see discussion by Udry et al. (2003)). If larger α is adopted, τ_ν becomes short compared with τ_{dep} , so that even relatively massive planets can migrate significantly, resulting in less clear deficit. We found that the two planet-depleted regions exist for other ΔZ_* as well, because the “planet desert” and the other are determined by the core accretion model for gas giants and $\tau_{\text{dep}}/\tau_\nu$, respectively. The observed distributions may exhibit the above two deficits and are consistent with our theoretical models with $\tau_{\text{dep}} \sim \tau_\nu$ at ~ 10 AU. Adopting the same condition, the simulated a -distribution of the gas giant planets can match with the observation [Figure 1c; also see Armitage et al. (2002) and Trilling et al. (2002)].

The 0.04 AU cut-off in the planets’ a distribution in our results may also be less abrupt and more consistent with the observed distributions if we adopt more relaxed inner boundary conditions. Many of giant planets may have migrated toward the vicinity of their host stars (Lin et al. 1996). The theoretically determined ratio of the giant planets with $a < 0.06\text{AU}$ (hot Jupiters) to those between 0.2–2AU is $\sim 1\text{--}10$ for stars with $Z_* \sim Z_\odot$, which is order of magnitude larger than its observed value (~ 0.2). This discrepancy suggests that more than 90% of the hot Jupiters which migrated to the stellar vicinity are either consumed (Sandquist et al. 1998) or tidally disrupted (Gu et al. 2003) by their host stars. Provided these disruptive events occur before the host stars become main sequence stars with relatively shallow convection zones, their apparent chemical homogeneity may be preserved to match the observation of the Pleiades cluster (Wilden et al. 2002). As the migrating protoplanets sweep planetesimals along their paths, this process may self-regulate the amount of residual heavy elements in the disk.

Figures 1c and d show that the normalized slopes of the a and M_p distributions are independent of ΔZ_* , because they are determined by $\tau_{\text{dep}}/\tau_\nu$ and the core accretion model for gas giants, respectively. In these figures, we do not include planets stalled at 0.04AU, since a large fraction of

them may plunge into their host stars.

Although the normalized distributions are independent of ΔZ_* , we found that formation and retention rates of gas giant planets rapidly increase with ΔZ_* . In order to account for the ΔZ_* -dependence in the M_p and a distribution, we first describe how planet formation depends on f_d [for details, see Paper I and Kokubo & Ida (2002)]. In a disk similar to the MMN ($f_d = f_{g,0} = 1$), for $a < a_{ice}$, the core growth is limited to $M_{c,iso} < M_\oplus$ where $M_\oplus$ is the Earth mass. Although growth beyond isolation (up to $\sim M_\oplus$) is possible for terrestrial planets, it is attainable only after severe gas depletion (Kominami & Ida 2002). At these regions, both $M_{c,iso}$ and $M_{c,noiso}$ are $< M_{c,acc}$ such that the cores cannot acquire a massive gaseous envelope. At the outer regions where $a \gg a_{ice}$, although $M_{c,iso} > M_{c,acc}$, the timescale required for M_c to reach $M_{c,acc}$ is $\gg \tau_{dep}$. The cores evolve into ice giants, similar to Uranus and Neptune. In the disk similar to the MMN, gas giants can form only slightly outside the ice boundary, because the cores may acquire $M_{c,acc}$ there within a few Myr and enter a rapid gas accretion phase prior to the global disk depletion. Gas giants can form inside a_{ice} in some massive disks (with $f_d \gtrsim 5$). Rocky cores are formed with $M_{c,iso} \lesssim M_\oplus$ in the inner regions and ice giants are formed with $M_c(\tau_{dep}) \lesssim M_\oplus$ in the outer regions of low-mass disks with $f_d \lesssim 0.5$, without formation of gas giants.

Around metal-deficient stars, formation efficiency of gas giants is low because disks with sufficiently large f_d are only at the tail end of the distribution. Around the metal-rich stars, the efficiency is high because both dM_c/dt and $M_{c,iso}$ increase rapidly with f_d (eqs. 3 and 4) and $M_{c,iso} \gtrsim M_{c,acc}$ even interior to a_{ice} . These account for the ΔZ_* -dependence. In order to directly compare with the observed frequency of extrasolar planets, we plot in Figures 2 the theoretical determination for fraction (η_J) of stars which bear giant planets detectable with the current Doppler survey, as a function of their metallicity. Here, the distribution of f_d is the same as that in Figures 1. For each f_d , a is selected as $\log(a_{j+1}/a_j) = 0.2$ ($j = 1, 2, \dots$). We do not include planets stalled at 0.04AU in the evaluation of η_J .

These results indicate η_J increases linearly with ΔZ_* , which is consistent with observational results of Fischer & Valenti (2003) (open circles in Figures 2). As shown in Figure 2a, η_J does not depend on the choice of τ_{KH} [$(b, c) = (9, 3), (10, 3)$ or $(11, 3.5)$; $\tau_{KH} = 10^b (M_p/M_\oplus)^{-c}$] nor that of asymptotic core mass [$\min(M_{c,iso}, M_{c,noiso})$ or $M_{c,noiso}$]. The reasons are as follows. As long as τ_{KH} has strong dependence on M_p , even significant change in b results in relatively small change of $M_{c,acc}$, the core mass that enables rapid gas accretion prior to disk gas depletion. Since in our simulations, most gas giant planets are formed with such parameter ranges where f_d and a are relatively large and $M_{c,iso} \gtrsim M_{c,noiso}$, the distributions hardly change even if $M_{c,noiso}$ is adopted in entire f_d and a (although it affects the distributions of terrestrial planets [Paper I]).

Although the absolute values of η_J are higher/lower for faster/slower core accretion, the dependence on ΔZ_* is similar (Figure 2b). If disk depletion is caused by viscous diffusion, the assumption that $\tau_{dep} \sim \tau_\nu$ at ~ 10 AU is reasonable. However, the observation of dense clusters suggest that gas depletion in protostellar disks is primarily driven by photo-evaporation (Johnstone et al. 1998)

in these clusters, where a few massive stars contribute to nearly the entire UV ionizing flux. In order to consider the possibility of rapid gas depletion in a dense cluster, we carried out the additional simulations with $\tau_{\text{dep}} = 10^5\text{--}10^6$ yrs. In this case, formation rates of giant planets are lower, resulting in smaller η_J (Figure 2b). But, the dependence on ΔZ_* does not change also in this case. This general agreement is a generic feature of the core accretion scenario and the assumption for f_d and $f_{g,0}$. It may not arise naturally from the gravitational instability scenario (Boss 1997).

4. Discussion

Through Monte Carlo simulation using the deterministic planet formation model developed in Paper I, we find that the formation probability of gas giant planets increases rapidly with the metallicity of their host stars. Our model quantitatively accounts for observational correlation between stellar metallicity and the fraction of stars harboring giant planets that are detectable with the current Doppler survey. Our model also explains the two observationally suggested planet-depleted regions in M_p - a distributions of extrasolar planets: the deficit in intermediate mass $M_p = 10\text{--}100M_\oplus$ and $a \lesssim 3\text{AU}$, and that in large mass $M_p \gtrsim 10^3M_\oplus$ and $a \lesssim 0.2\text{AU}$.

Our model can be tested by more detailed observation. We show that the normalized slope of the a and M_p distributions of gas giants and the planet-depleted regions are independent of ΔZ_* , because these are regulated by $\tau_{\text{dep}}/\tau_\nu$ and the core accretion model. The lower limit mass $M_{p,\text{lim}}$ of hot Jupiters depend on ΔZ_* as $M_{p,\text{lim}} \propto 10^{-(3/2)\Delta Z_*}$ (Paper I). The outer boundary radius of the distribution of gas giants $a_{\text{out}} \propto 10^{(10/27)\Delta Z_*}$ (Paper I), although interactions among gas giants, which are not taken into account in the present paper, may make the boundary less clear (Marzari & Weidenschilling 2002). These predictions of ΔZ_* -dependence can be tested by statistics of extrasolar planets with increased detection number in a few years.

Our results provide supporting evidences for the core-accretion scenario and they suggest that the detection of gas giant planets may be used to infer the presence of rocky planets around the same host stars. "Habitable planets" which can retain liquid water on their surface (Kasting et al. 1993) may have $0.1M_\oplus \lesssim M_p \lesssim 10M_\oplus$ and $a \sim 0.8\text{--}1.5 \times (M_*/M_\odot)^2 \text{AU}$. The results in Figure 1 indicate that the frequency of "habitable planets" is comparable to or more than that of detectable giant planets (at present, more than 100 planets have been detected around F, G, K dwarfs within 50 pc), although some fraction of the habitable planets may be destabilized by migrating giant planets.

REFERENCES

- Alibert, Y., Mordasini, C., Benz, W. 2004, A&A, 417, L25
- Aarseth, S. J., Lin, D. N. C., Palmer, P. L. 1993, ApJ, 403,351
- Armitage, P. J., Livio, M., Lubow, S. H., & Pringle, J. E. 2002, MNRAS, 334, 248
- Artymowicz, P. 1993, ApJ, 419, 166
- Beckwith, S. V. W. & Sargent, A. I., 1996, Nature, 383, 139
- Bodenheimer, P., & Pollack, J. B. 1986, Icarus, 67, 391
- Boss, A. P. 1997, Science, 276, 1836
- Calvet, N., Hartmann, L. & Strom, S. E. 2000, in Protostars and Planets IV, ed. V. Mannings, A. P. Boss and S. S. Russell (Tucson:Univ. of Arizona Press), 377
- Desidera, S. et al. 2004, A&A, in press
- Fischer, D. A. & Valenti, J. A. 2003. in Scientific Frontiers in Research on Extrasolar Planets, ASP Conference Series 294, ed. D. Deming & S. Seager. (San Francisco: ASP) , 117.
- Goldreich, P., & Tremaine, S. 1980, ApJ, 241, 425
- Greenberg, R., Hartmann, W. K., Chapman, C. R., & Wacker, J. F. 1978, Icarus 35, 1
- Gu, P., Lin, D. N. C., & Bodenheimer, P. H. 2003, ApJ, 588, 509
- Haisch, K. E., Lada, E. A. & Lada, C. J. 2001, ApJ, 553, L153
- Hayashi, C. 1981, Prog. Theor. Phys. Suppl., 70, 35
- Ida, S. & Lin, D. N. C. 2004, ApJ, 604, 388 (Paper I)
- Ida, S., Bryden, G., Lin, D. N. C., & Tanaka, H. 2000, ApJ, 534, 428
- Ikoma, M., Nakazawa, K. & Emori, E. 2000, ApJ, 537, 1013
- Inaba, S., Wetherill, G. & Ikoma, M. 2003, Icarus, 410, 711
- Johnstone, D., Hollenbach, D. & Bally, J. 1998, ApJ, 499, 758
- Kasting, J. F., Whitmire, D. P. & Reynolds, R. T. 1993, Icarus,101, 108
- Kokubo, E. & Ida, S. 1996, Icarus, 123, 180
- . 1998, Icarus, 131, 171

- . 2002, *ApJ*, 581, 666
- Kominami, J. & Ida, S. 2002, *Icarus*, 157, 43
- Lin, D. N. C., Bodenheimer, P. & Richardson, D. 1996, *Nature*, 380, 606
- Lin, D. N. C. & Papaloizou, J. C. B. 1985, *Protostars and Planets II*, ed. D. C. Black & M. S. Matthews (Tucson: Univ. of Arizona Press), 981
- Lin, D. N. C., & Papaloizou, J. C. B. 1993, in *Protostars and Planets III*, ed. E. H. Levy and J. I. Lunine (Tucson: Univ. of Arizona Press), 749
- Lubow, S. H., Seibert, M. & Artymowicz, P. 1999, *ApJ*, 526, 1001
- Nelson, R. P. & Papaloizou, J. C. B. 2004, *MNRAS*, 350, 849
- Marzari, F. & Weidenschilling, S. 2002, *Icarus*, 156, 570
- Mizuno, H. 1980, *Prog. Theor. Phys. Suppl.*, 64, 54
- Murray, N. & Chaboyer, B. 2001, *ApJ*, 566, 442
- Murray, N., Hansen, B., Holman, M., & Tremaine, S. 1998, *Science*, 279, 69
- Pollack, J. B., Hollenbach, D., Beckwith, S., Simonelli, D. P., Roush, T., & Fong, W. 1994, *ApJ*, 421, 615
- Pollack, J. B., Hubickyj, O., Bodenheimer, P., Lissauer, J. J., Podolak, M., & Greenzweig, Y. 1996, *Icarus*, 124, 62
- Quillen, A. C. 2002, *AJ*, 124, 400
- Rafikov, R. R. 2003, *ApJ*, 125, 922
- Rice, W. K. M. & Armitage, P. J. 2003, *ApJ*, 598, L55
- Sandquist, E., Taam, R. E., Lin, D. N. C. & Burkert, A. 1998, *ApJ*, 506, L65
- Santos, N. C., Israelian, G. & Mayor, M. 2004, *A&A*, 415, 1153
- Shakura, N. I. & Sunyaev, R. A. 1973, *A&A*, 24, 337
- Tanaka, H. & Ida, S. 1999, *Icarus*, 125, 302
- Tanaka, H. & Ida, S. 1999, *Icarus*, 139, 350
- Trilling, D. E., Lunine, J. I. & Benz, W. 2002, *ã*, 394, 241
- Udry, S., Mayor, M., & Santos, N. C. 2003, *ã*, 407, 369

Ward, W. 1986, *Icarus*, 67, 164

———. 1993, *Icarus*, 106, 274

———. 1997, *Icarus*, 126, 261

Wetherill, G. W. & Stewart, G. R. 1989, *Icarus*, 77, 330

Wilden, B. S., Jones, B. F., Lin, D. N. C. & Soderblom, D. R. 2002. *AJ*, 124, 2799

ACKNOWLEDGMENTS. We thank Debra Fischer, Greg Laughlin and Masahiro Ikoma for useful discussions and the anonymous referee for helpful comments. This work is supported by the NASA Origins program, JPL SIM program, NSF and JSPS.

CORRESPONDENCE should be addressed to S. I. (ida@geo.titech.ac.jp).

FIGURE LEGENDS

Fig.1

Theoretically predicted mass (M_p) and semi-major axis (a) distributions of extrasolar planets. For parameters, see text. In panel (a), the computed result with $\Delta Z_* = 0$ is shown. Totally 20000 planets are calculated. For comparison, observational data are plotted in panel (b). (Since orbital inclinations are unknown, minimum masses are plotted). The observational limits in the current Doppler survey are marked by dotted lines. One dimensional normalized distributions of observable planets is plotted as a function of a (panel c) and M_p (panel d). Large open circles are observed results. Small filled circles, pentagons, squares, triangles, and crosses express the theoretical results with $\Delta Z_* = 0.5, 0.25, 0, -0.25$ and -0.5 , respectively. Here 1000 observable planets are used in each ΔZ_* result.

Fig.2

The fraction (η_J) of stars which bears giant planets currently detectable with Doppler survey, as a function of their metallicity (ΔZ_*). Large open circles are observational results (Fischer & Valenti 2003). Other small symbols with lines represent theoretical predictions. Filled circles express the nominal case where core accretion is given by Eq. (3) with $m = 10^{22}$ g, the asymptotic masses are $\min(M_{c,iso}, M_{c,noiso})$, and $\tau_{KH} = 10^b (M_p/M_\oplus)^{-c}$ yrs with $(b, c) = (10, 3)$. In panel (a), crosses, filled triangles and squares express the cases with $M_{c,noiso}$, $(b, c) = (9, 3)$, and $(b, c) = (11, 3.5)$, respectively. In panel (b), crosses, filled triangles and squares express the cases with 3 times faster core accretion rate, 3 times slower one, and $\tau_{dep} = 10^5$ – 10^6 yrs, respectively.




Fig. 1.—




Fig. 2.—